PATENT COOPERATION TREATY PCT

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY

(Chapter II of the Patent Cooperation Treaty)

(PCT Article 36 and Rule 70)

	, 				
Applicant's or agent's file reference 42968PCX329/29 KM	FOR FURTHER ACTION	See Form PCT/IPEA/416			
International application No. PCT/NZ2004/000184	International filing date (day/month/yell 13 August 2004	Priority date (day/month/year) 15 August 2003			
International Patent Classification (IPC) or					
Applicant AGRESEARCH LIMITED et al					
This report is the international prelimin Authority under Article 35 and transmit	ary examination report, established by the ted to the applicant according to Article				
2. This REPORT consists of a total of 4	sheets, including this cover sheet.	·			
3. This report is also accompanied by AN	NEXES, comprising:				
a. X (sent to the applicant and to the	e International Bureau) a total of 17 s	heets, as follows:			
		amended and are the basis for this report and/or			
sheets containing rectification Administrative Instruction	ations authorized by this Authority (see last).	Rule 70.16 and Section 607 of the			
	,	nsiders contain an amendment that goes beyond			
the disclosure in the international application as filed, as indicated in item 4 of Box No. I and the Supplemental Box.					
b. (sent to the International Burea	au only) a total of (indicate type and nun	nber of electronic carrier(s)) , containing			
a sequence listing and/or table	related thereto, in computer readable for see Section 802 of the Administrative In	m only, as indicated in the Supplemental Box			
4. This report contains indications relating		su ucuons).			
X Box No. I Basis of the repo	rt .	·			
Box No. II Priority					
Box No. III Non-establishme	nt of opinion with regard to novelty, inv	rentive step and industrial applicability			
Box No. IV Lack of unity of	invention				
Box No. V Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement					
Box No. VI Certain documen					
Box No. VII Certain defects in	Box No. VII Certain defects in the international application				
Box No. VIII Certain observations on the international application					
Date of submission of the demand Date of completion of the report					
1 February 2005	26 July 2005				
Name and mailing address of the IPEA/AU	Authorized Offic	er			
AUSTRALIAN PATENT OFFICE					
PO BOX 200, WODEN ACT 2606, AUSTRA E-mail address: pct@ipaustralia.gov.au	S. Chew				
Facsimile No. (02) 6285 3929	(02) 6283 2248				

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY

International application No.

PCT/NZ2004/000184

Box	Box No. I Basis of the report				
1.	With regard to the language, this report is based on the international application in the language in which it was filed, unless otherwise indicated under this item.				
•	This report is based on translations from the original language into the following language, which is the language of a translation furnished for the purposes of:				
	international search (under Rules 12.3 and 23.1 (b))				
	publication of the international application (under Rule 12.4)				
	international preliminary examination (under Rules 55.2 and/or 55.3)				
2.	With regard to the elements of the international application, this report is based on (replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report):				
	the international application as originally filed/furnished				
	X the description:				
1	pages 1-33 as originally filed/furnished				
	pages* received by this Authority on with the letter of pages* received by this Authority on with the letter of				
	X the claims:				
	pages as originally filed/furnished				
	pages* as amended (together with any statement) under Article 19	_			
	pages*34-50 received by this Authority on 1 February 2005 with the letter of 1 February 2005 pages* received by this Authority on with the letter of	,			
	X the drawings:				
	pages 1/5 - 5/5 as originally filed/furnished				
	pages* received by this Authority on with the letter of				
	pages* received by this Authority on with the letter of				
	a sequence listing and/or any related table(s) - see Supplemental Box Relating to Sequence Listing.				
3.	The amendments have resulted in the cancellation of:				
	the description, pages				
	the claims, Nos.				
	the drawings, sheets/figs				
	the sequence listing (specify):				
	any table(s) related to the sequence listing (specify):				
4.	This report has been established as if (some of) the amendments annexed to this report and listed below had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).				
	the description, pages				
	the claims, Nos.				
	the drawings, sheets/figs				
	the sequence listing (specify):				
	any table(s) related to the sequence listing (specify):				
*	* If item 4 applies, some or all of those sheets may be marked "superseded."				

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY

International application No.

PCT/NZ2004/000184

Box No. V	Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability;
citation	s and explanations supporting such statement

Statement		
Novelty (N)	Claims 1-49	YES
	Claims	NO
Inventive step (IS)	Claims 1-49	YES
	Claims	NO
Industrial applicability (IA)	Claims 1-49	YES
	Claims	NO

2. Citations and explanations (Rule 70.7)

This report has considered the following documents cited in the International Search Report:

- D1 WO 2003/105868
- D2 Miles C. et al.
- D3 Munday-Finch S. et al. J. Agric. Food Chem. 1995
- D4 Munday-Finch S. et al. J. Agric. Food Chem. 1998
- D5 Munday-Finch S. et al. J. Agric. Food Chem. 1997
- D6 Derwent Abstract Accession No. 92-308267/38
- D7 Munday-Finch S. et al. J. Agric. Food Chem. 1996

NOVELTY (N), INVENTIVE STEP (IS): Claims 1-49

D1 discloses lolitrems A, B, C, E, F, H, N, lolitrem N-31-epimer, lolitriol, lolilline, lolitriol, lolicines A and B and their use as potassium channel blockers for the treatment of ocular hypertension or glaucoma (see pages 5, 7, 13 and claim 1).

D2 discloses the isolation and structures of lolitrems B and E including their biosynthetic route from lolitriol (see abstract and figure 1).

D3 discloses the isolation of lolitrem A, its structure and structures of lolitrems B, C and E (see abstract and figure 1).

D4 discloses the isolation of lolicines A and B, lolitriol and lolitrem N and has provided evidence for 31-epilolitrem N and 31-epilolitrem F (see abstract and figure 1).

D5 discloses lolilline, lolitrems A, B, E and lolitriol (see figures 1 and 3).

D6 discloses some lolitrem derivatives used for the preparation of haptens for the production of antibodies.

D7 discloses lolitrem F, lolitrem B, 31-epilolitrem B, 31-epilolitrem F and lolitriol (see abstract, figures 1 and 4).

INTERNATIONAL PRELIMINARY REPORT ON PATENTABILITY

International application No.

PCT/NZ2004/000184

Supplem	ental	Box
---------	-------	-----

case the space in any of the preceding boxes is not sufficient.

Continuation of Box No. V:

None of D1-D7 disclose or fairly suggest alone or in combination, a method of preventing repolarisation or hyperpolarisation of a cell wherein the cell contains a BK channel, comprising the administration to the cell of a composition containing a BK channel antagonist as defined in the claims, or a composition comprising a BK channel antagonist compound containing the moiety shown in structures (VII), (IX), (XII) and (XIII).

Therefore claims 1-49 are novel and have an inventive step.

INDUSTRIAL APPLICABILITY (IA): Claims 1-49

Claims 1-49 have industrial applicability.

34

IAP20 Racid PCT/FTO 17 JAN 2006

WHAT WE CLAIM IS:

5

10

15

1. A method of preventing repolarisation or hyperpolarisation of a cell, wherein the cell contains a BK channel, including the administration to the cell of at least one pharmacologically effective amount of composition containing a BK channel antagonist containing the moiety shown in structure (I):

STRUCTURE (I)

or derivatives thereof.

- 2. The method as claimed in claim 1 wherein the derivatives of structure (I) are selected from the group consisting of: salts, analogues, isomers, and combinations thereof.
 - 3. The method as claimed in claim 1 or claim 2 wherein the antagonist compound is selected from the group consisting of: lolitrem B, lolitrem A, lolitrem F, 31-*epi*lolitrem F, 31-*epi*lolitrem B, lolitrem E, lolitrem E acetate, lolitrem L, lolitrem G, lolitrem C, lolitrem M, lolitriol, lolitriol acetate, lolitrem N, lolitrem J, lolitrem H, lolitrem K, lolicine A and B, 30-desoxy lolitrem B-30α-ol, 30-desoxy-31-*epi*lolitrem B-30α-ol, 30-desoxylolitrem B-30-ene lolilline and combinations thereof.
 - 4. The method as claimed in claim 1 or claim 2 wherein the antagonist compound is selected from the group consisting of:

STRUCTURE (II)

which includes compounds selected from the group consisting of: lolitrem B = 31α , 35β stereochemistry; 31-epilolitrem B = 31β , 35β stereochemistry; lolitrem F = 31α , 35α ; 31-epilolitrem F = 31β , 35α ;

STRUCTURE (III)

which includes compounds selected from the group consisting of: lolitrem E = 31α , 35β stereochemistry where R = H or acetate; lolitrem L = 31α , 35α

stereochemistry where R = H or acetate;

STRUCTURE (IV)

which includes compounds selected from the group consisting of: lolitrem A = 31α , 35β stereochemistry; lolitrem G = 31α , 35α stereochemistry;

STRUCTURE (V)

which includes compounds selected from the group consisting of: lolitriol; = 31α , 35β stereochemistry where R₁ = H or acetate and R₂ = H; lolitrem N = 31α , 35α stereochemistry where R₁=H or acetate and R₂=H; Lolitrem J = 31α , 35β stereochemistry where R₁ = H or acetate and R₂ = acetate;

STRUCTURE (VI)

10

which includes lolitrem H = 31α , 35β stereochemistry where R = H or acetate;

37 STRUCTURE (VII)

which includes lolitrem K = 31α , 35β stereochemistry, where R = H or acetate;

STRUCTURE (VIII)

which includes lolilline = 31α , 35β stereochemistry;

STRUCTURE (IX)

which includes lolitrem M = 31α , 35β stereochemistry;

STRUCTURE (X)

10

which includes lolicine A = 31α , 35β stereochemistry;

STRUCTURE (XI)

which includes lolicine B = 31α , 35β stereochemistry;

STRUCTURE (XII)

which includes compounds selected from the group consisting of: 30-desoxylolitrem B-30 α -ol = 31 α , 35 β stereochemistry; 30-desoxy-31-*epi*lolitrem B-30 α -ol = 31 β , 35 β stereochemistry;

STRUCTURE (XIII)

10

which includes 30-desoxylolitrem B-30-ene = 35β stereochemistry;

and combinations of the above compounds.

- 4. The method as claimed in any of the above claims wherein the composition further includes pharmaceutically and physiologically acceptable carriers.
- 5 5. The method as claimed in claim 4 wherein the pharmaceutically and physiologically acceptable carriers include components selected from the group including; fillers; excipients; modifiers; humectants; stabilisers; emulsifiers; diluents; and other formulation components such as a use of a lipid vehicle.
- 6. The method as claimed in any of the above claims wherein the composition is
 administered in a form selected from the group including: an injection; a tablet; a
 capsule; a suppository; an injection; a suspension; a drink or tonic; a syrup; a
 powder; an ingredient in solid or liquid foods; a nasal spray; a sublingual wafer; a
 transdermal patch; a transdermal injection; and combinations thereof.
- 7. The method as claimed in any of the above claims wherein the BK channel
 antagonist compound or compounds are extracted from endophyte-infected plants and seeds.
 - 8. The method as claimed in any of claims 1 to 6 wherein the BK channel antagonist compound or compounds are extracted from fungal cultures.
 - 9. The method as claimed in any of claims 1 to 6 wherein the BK channel antagonist compound or compounds are derived by chemical synthesis.

- 10. The method as claimed in any of claims 1 to 6 wherein the BK channel antagonist compound or compounds are extracted from heterologous expression systems including but not limited to bacteria, yeast, fungi, plants and animal cells.
- 11. The method as claimed in claim 7 wherein the perennial ryegrass seed is from

Lolium perenne.

- 12. The method as claimed in any of the above claims wherein the BK channel antagonist compound or compounds has activity against both alpha (α) subunit and alpha plus beta (β) accessory subunit (β_1 to β_4) channels.
- 13. The method as claimed in any of claims 1 to 4 wherein, for lolitrem B, the degree of antagonist inhibition is approximately 97% for a composition containing approximately 20nM lolitrem B.
 - 14. The method as claimed in any of claims 1 to 4 wherein, for lolitrem B, the half maximal degree of antagonist inhibition (IC_{50}) is found for a composition containing approximately 3.7 \pm 0.4 nM of lolitrem B.
 - 15. The method as claimed in any of claims 1 to 4 wherein, for lolitriol, the degree of antagonist inhibition is approximately 100% for a composition containing approximately 1000 nM lolitriol.
- 16. The method as claimed in any of claims 1 to 4 wherein, for lolitriol, the half
 maximal degree of antagonist inhibition (IC₅₀) is found for a composition containing approximately 195 nM of lolitriol to inhibit α and β₁ BK channel activity
 - 17. The method as claimed in any of claims 1 to 4 wherein, for lolitriol, the half maximal degree of antagonist inhibition (IC₅₀) is found for a composition containing approximately 536 \pm 16 nM of lolitriol to inhibit α and β_4 activity.
- 18. The method as claimed in any of claims 1 to 4 wherein, for 31-epilolitrem B, the degree of antagonist inhibition is approximately 100% for a composition containing approximately 200nM 31-epilolitrem B.
 - 19. The method as claimed in any of claims 1 to 4 wherein, for 31-epilolitrem B, the half maximal degree of antagonist inhibition (IC₅₀) is found for a composition

containing approximately 58 ± 6 nM of 31-epilolitrem B to inhibit α and β_1 activity.

- 20. The method as claimed in any of claims 1 to 4 wherein, for 31-epilolitrem B, the half maximal degree of antagonist inhibition (IC₅₀) is found for a composition containing approximately 49 nM of 31-epilolitrem B to inhibit α and β_4 activity.
- 21. The method as claimed in any of claims 1 to 4 wherein, for lolitrem E, the degree of antagonist inhibition is approximately 100% for a composition containing approximately 100 nM lolitrem E.
 - 22. The method as claimed in any of claims 1 to 4 wherein the antagonist effect of the composition is not able to be reversed by wash out for concentrations of 10 nM or greater of lolitrem B compound.
 - 23. Use of a composition for preventing repolarisation or hyperpolarisation of a cell that contains a BK channel wherein a pharmacologically effective amount of the composition is administered to the cell and wherein the composition contains at least one BK channel antagonist of the moiety shown in structure (I):

15

20

10

STRUCTURE (I)

or derivatives thereof.

24. The use as claimed in claim 23 wherein the derivatives of structure (I) are selected from the group consisting of: salts, analogues, isomers, and combinations thereof.

- 25. The use as claimed in claim 23 or claim 24 wherein the antagonist compound is selected from the group consisting of: lolitrem B, lolitrem A, lolitrem F, 31
 epilolitrem F, 31-epilolitrem B, lolitrem E, lolitrem E acetate, lolitrem L, lolitrem G, lolitrem C, lolitrem M, lolitriol, lolitriol acetate, lolitrem N, lolitrem J, lolitrem H, lolitrem K, lolicine A and B, 30-desoxy lolitrem B-30α-ol, 30-desoxy-31-epilolitrem B-30α-ol, 30-desoxylolitrem B-30-ene lolilline and combinations thereof.
 - 26. The use as claimed in claim 23 or claim 24 wherein the antagonist compound is selected from the group consisting of:

•

STRUCTURE (II)

which includes compounds selected from the group consisting of: lolitrem B = 31α , 35β stereochemistry; 31-*epi*lolitrem B = 31β , 35β stereochemistry; lolitrem F = 31α , 35α ; 31-*epi*lolitrem F = 31β , 35α ;

15

10

5

STRUCTURE (III)

which includes compounds selected from the group consisting of: lolitrem E = 31α , 35β stereochemistry where R = H or acetate; lolitrem L = 31α , 35α stereochemistry where R = H or acetate;

STRUCTURE (IV)

which includes compounds selected from the group consisting of: lolitrem A = 31α , 35β stereochemistry; lolitrem G = 31α , 35α stereochemistry;

STRUCTURE (V)

5

10

which includes compounds selected from the group consisting of: lolitriol; = 31α , 35β stereochemistry where R₁ = H or acetate and R₂ = H; lolitrem N = 31α , 35α stereochemistry where R₁=H or acetate and R₂=H; Lolitrem J = 31α , 35β stereochemistry where R₁ = H or acetate and R₂ = acetate;

STRUCTURE (VI)

which includes lolitrem H = 31α , 35β stereochemistry where R = H or acetate;

Amended Sheet IPEA/AU

STRUCTURE (VII)

which includes lolitrem K = 31α , 35β stereochemistry, where R = H or acetate;

5

STRUCTURE (VIII)

which includes Iolilline = 31α , 35β stereochemistry;

STRUCTURE (IX)

10

which includes lolitrem $M = 31\alpha$, 35β stereochemistry;

STRUCTURE (X)

which includes lolicine $A = 31\alpha$, 35β stereochemistry;

STRUCTURE (XI)

5

which includes lolicine B = 31α , 35β stereochemistry;

STRUCTURE (XII)

which includes compounds selected from the group consisting of: 30-10 desoxylolitrem B-30 α -ol = 31 α , 35 β stereochemistry; 30-desoxy-31-*epi*lolitrem B-30 α -ol = 31 β , 35 β stereochemistry;

STRUCTURE (XIII)

which includes 30-desoxylolitrem B-30-ene = 35β stereochemistry;

and combinations of the above compounds.

10

- 5 27. The use as claimed in any of the above claims wherein the composition further includes pharmaceutically and physiologically acceptable carriers.
 - 28. The use as claimed in claim 27 wherein the pharmaceutically and physiologically acceptable carriers include components selected from the group including; fillers; excipients; modifiers; humectants; stabilisers; emulsifiers; diluents; and other formulation components such as a use of a lipid vehicle.
 - 29. The use as claimed in any of claims 23 to 28 wherein the composition is administered in a form selected from the group including: an injection; a tablet; a capsule; a suppository; an injection; a suspension; a drink or tonic; a syrup; a powder; an ingredient in solid or liquid foods; a nasal spray; a sublingual wafer; a transdermal patch; a transdermal injection; and combinations thereof.
 - 30. The use as claimed in any of claims 23 to 29 wherein the BK channel antagonist compound or compounds are extracted from endophyte-infected plants and seeds.
- 31. The use as claimed in any of claims 23 to 29 wherein the BK channelantagonist compound or compounds are extracted from fungal cultures.

32. The use as claimed in any of claims 23 to 29 wherein the BK channel antagonist compound or compounds are derived by chemical synthesis.

5

- 33. The use as claimed in any of claims 23 to 29 wherein the BK channel antagonist compound or compounds are extracted from heterologous expression systems including but not limited to bacteria, yeast, fungi, plants and animal cells.
- 34. The use as claimed in claim 30 wherein the perennial ryegrass seed is from *Lolium perenne*.
- 35. The use as claimed in any of claims 23 to 34 wherein the BK channel antagonist compound or compounds has activity against both alpha (α) subunit and alpha plus beta (β) accessory subunit (β_1 to β_4) channels.
- 36. The use as claimed in any of claims 23 to 26 wherein, for lolitrem B, the degree of antagonist inhibition is approximately 97% for a composition containing approximately 20nM lolitrem B.
- 37. The use as claimed in any of claims 23 to 26 wherein, for lolitrem B, the half maximal degree of antagonist inhibition (IC_{50}) is found for a composition containing approximately 3.7 \pm 0.4 nM of lolitrem B.
 - 38. The use as claimed in any of claims 23 to 26 wherein, for lolitriol, the degree of antagonist inhibition is approximately 100% for a composition containing approximately 1000 nM lolitriol.
- 39. The use as claimed in any of claims 23 to 26 wherein, for lolitriol, the half maximal degree of antagonist inhibition (IC₅₀) is found for a composition containing approximately 195 nM of lolitriol to inhibit α and β_1 BK channel activity
 - 40. The use as claimed in any of claims 23 to 26 wherein, for lolitriol, the half maximal degree of antagonist inhibition (IC_{50}) is found for a composition containing

approximately 536 ±16 nM of lolitriol to inhibit α and β_4 activity.

- 41. The use as claimed in any of claims 23 to 26 wherein, for 31-epilolitrem B, the degree of antagonist inhibition is approximately 100% for a composition containing approximately 200nM 31-epilolitrem B.
- 42. The use as claimed in any of claims 23 to 26 wherein, for 31-epilolitrem B, the half maximal degree of antagonist inhibition (IC₅₀) is found for a composition containing approximately 58 ±6 nM of 31-epilolitrem B to inhibit α and β₁ activity.
 - 43. The use as claimed in any of claims 23 to 26 wherein, for 31-epilolitrem B, the half maximal degree of antagonist inhibition (IC₅₀) is found for a composition containing approximately 49 nM of 31-epilolitrem B to inhibit α and β_4 activity.
 - 44. The use as claimed in any of claims 23 to 26 wherein, for lolitrem E, the degree of antagonist inhibition is approximately 100% for a composition containing approximately 100 nM lolitrem E.
- 45. The use as claimed in any of claims 23 to 26 wherein the antagonist effect of
 the composition is not able to be reversed by wash out for concentrations of 10 nM or greater of lolitrem B compound.
 - 46. A composition that contains a pharmacologically effective amount of at least one BK channel antagonist compound containing the moiety shown in structure (VII):

20

10

STRUCTURE (VII)

which includes lolitrem $K = 31\alpha$, 35β stereochemistry, where R = H or acetate.

47. A composition that contains a pharmacologically effective amount of at least one BK channel antagonist compound containing the moiety shown in structure (IX):

STRUCTURE (IX).

which includes lolitrem M = 31α , 35β stereochemistry.

5

10

48. A composition that contains a pharmacologically effective amount of at least one BK channel antagonist compound containing the moiety shown in structure (XII):

STRUCTURE (XII)

which includes compounds selected from the group consisting of: 30-desoxylolitrem B-30 α -ol = 31 α , 35 β stereochemistry; 30-desoxy-31-*epi*lolitrem B-30 α -ol = 31 β , 35 β stereochemistry.

49. A composition that contains a pharmacologically effective amount of at least one BK channel antagonist compound wherein the antagonist compound is structure (XIII):

STRUCTURE (XIII)

which includes 30-desoxylolitrem B-30-ene = 35β stereochemistry.